

ON REDUCED Q -FUNCTIONS

Tatsuhiro NAKAJIMA

*Department of Physics, Tokyo Metropolitan University
Minami-Ohsawa 1-1, Hachioji-shi, Tokyo 192-03, Japan*

and

Hiro-Fumi YAMADA

*Department of Mathematics, Tokyo Metropolitan University
Minami-Ohsawa 1-1, Hachioji-shi, Tokyo 192-03, Japan*

To Kiyosato Okamoto with affection and admiration

Abstract

Schur's Q -functions with reduced variables are discussed by employing a combinatorics of strict partitions. They are called reduced Q -functions. We give a description of the linear relations among reduced Q -functions.

§0 Introduction

Q -functions were introduced by Schur in his study of projective representations of symmetric groups. They are symmetric functions and, if we express them in terms of the power sum symmetric functions, each coefficient essentially gives the spin character of the symmetric group.

This note deals with the “ r -reduced” Q -functions which are defined by putting $p_{jr} = 0$ for $j = 1, 3, 5, \dots$ in the power sum expression. When $r = p$ is a prime number, they play a role in p -modular projective representations of symmetric groups.

In a previous work [NY] we showed that r -reduced Q -functions are weight vectors of the basic representation of the affine Lie algebra $A_{2t}^{(2)}$ ($r = 2t + 1$) and chose a proper basis for each weight space.

To be more precise, let α_i (resp. $\alpha_i^\vee$) ($0 \leq i \leq t$) be the simple roots (resp. coroots) of the affine Lie algebra $A_{2t}^{(2)}$ and $\delta = 2 \sum_{i=0}^{t-1} \alpha_i + \alpha_t$ be the fundamental imaginary root. The irreducible representation with highest weight Λ_0 is called the basic representation, where $\Lambda_0(\alpha_i^\vee) = \delta_{i0}$. The set of weights is described by

$$P = \{W\Lambda_0 - n\delta ; n \in \mathbb{N}\},$$

where W is the Weyl group. This basic representation can be realized on the polynomial ring $\mathbb{C}[t_j ; j \geq 1, \text{ odd and } j \not\equiv 0 \pmod{r}]$. In this realization each r -reduced Q -function turns out to be a weight vector. We answered in [NY] the question to which weight the given r -reduced Q -function belongs, by using a combinatorics of strict partitions, i.e., bar cores and bar quotients.

By virtue of the above result we further investigate the r -reduced Q -functions themselves. The result in this note is an explicit description of the linear relations satisfied by r -reduced Q -functions. We remark that such a description has been obtained for “ r -reduced Schur functions” [ANY1,2].

§1 Combinatorics of strict partitions

We first present a collection of definitions and results concerning strict partitions and Schur's Q -functions, mainly referring to the new book of Macdonald [M].

The set of all partitions of n is denoted by $\mathcal{P}_n$ and the set of all partitions by $\mathcal{P}$. A partition $\lambda = (\lambda_1, \dots, \lambda_l)$ is said to be strict if $\lambda_1 > \lambda_2 > \dots > \lambda_l > 0$. The set of all strict partitions of n is denoted by $\mathcal{SP}_n$ and the set of all strict partitions by $\mathcal{SP}$.

To each element $\lambda \in \mathcal{SP}$ we can associate the shifted diagram $S(\lambda)$, which is obtained from the usual Young diagram of λ by shifting the i -th row $(i-1)$ -positions to the right. The j -th square in the i -th row will be called the (i, j) -square. For i, j and k with $k \geq i$, we define an $(i, j)_k$ -bar of λ as the set of squares $(a, b) \in S(\lambda)$ satisfying (1) if $k > i$, then $a = i$ or k , (2) if $k = i$, then $a = i$, $b \geq j$ and $j-1 \neq \lambda_{a'}$ for $a' \geq i+1$. The length of a bar is a number of squares it contains. A bar of length r will be often called an r -bar. Throughout this note we always assume that r is an odd number.

A strict partition is an r -bar core if it contains no r -bars. Given a strict partition λ we obtain another strict partition by removing an r -bar and rearranging the rows in descending order. By repeating this process as often as possible we shall end up with an r -bar core λ^c , which is called the r -bar core of λ .

For $0 \leq i \leq r-1$ we define sequences

$$X^i(\lambda) = \{x \in \mathbb{N}; \lambda_k = rx + i \text{ for some } k\}.$$

The r -bar quotient $\lambda^q = (\lambda^0, \lambda^1, \dots, \lambda^t)$ ($t = (r-1)/2$) is a $(t+1)$ -tuple of partitions given by $\lambda^0 = X^0(\lambda)$ and $\lambda^i = (X^i(\lambda) | X^{r-i}(\lambda))$ ($1 \leq i \leq t$) in Frobenius notation. Note that λ^0 is a strict partition. We remark that any r -bar core and r -bar quotient uniquely determines a strict partition [O]. In view of this one-to-one correspondence we often write (λ^c, λ^q) instead of λ , e.g., $Q_{(\lambda^c, \lambda^q)}(x)$, etc.

We introduce r -bar sign for strict partitions. Let $\bar{H}$ be the $(i, j)_k$ -bar of length r in

$\lambda \in \mathcal{SP}$. The leglength of $\bar{H}$ is defined by

$$b(\bar{H}) = \begin{cases} \lambda_k + \#\{a; \lambda_i > \lambda_a > \lambda_k\} & \text{if } k > i \\ \#\{a; \lambda_i > \lambda_a > \lambda_i - r\} & \text{if } k = i. \end{cases}$$

If the strict partition μ is obtained from λ by removing r -bars $H_1, H_2, \dots, H_q$ successively, then the r -bar sign of λ relative to μ is defined by

$$\delta_{\bar{r}}(\lambda, \mu) = \prod_{m=1}^q (-1)^{b(\bar{H}_m)}.$$

We set $\delta_{\bar{r}}(\lambda) = \delta_{\bar{r}}(\lambda, \lambda^c)$ and call it the r -bar sign of λ . We should remark that $\delta_{\bar{r}}(\lambda, \mu)$ does not depend on the choice of r -bars being removed in going from λ to μ [MO].

Let $\Lambda_{\mathbb{Q}}$ denote the graded $\mathbb{Q}$ -algebra of symmetric functions in countably many independent variables $x = (x_1, x_2, \dots)$. Among various bases of $\Lambda_{\mathbb{Q}}$, we just mention here that the Schur functions $s_{\lambda}(x)$ ($\lambda \in P$) are convenient for our point of view. Let $\Gamma_{\mathbb{Q}}$ be the subalgebra of $\Lambda_{\mathbb{Q}}$ generated by $1, p_1, p_3, p_5, \dots$;

$$\Gamma_{\mathbb{Q}} = \mathbb{Q}[p_1, p_3, p_5, \dots],$$

where $p_j = x_1^j + x_2^j + \dots$ is the j -th power sum.

Schur's Q -functions $Q_{\lambda}(x)$ ($\lambda \in \mathcal{P}$) are defined by specializing to $t = -1$ in the Hall-Littlewood symmetric functions $Q_{\lambda}(x; t)$. This specialization implies that $Q_{\lambda}(x) = 0$ unless $\lambda \in \mathcal{SP}$. It is known that $\{Q_{\lambda}(x); \lambda \in \mathcal{SP}\}$ gives a basis of $\Gamma_{\mathbb{Q}}$. When $\mu = \sigma\lambda = (\lambda_{\sigma(1)}, \dots, \lambda_{\sigma(l)})$ ($\sigma \in \mathfrak{S}_l$) is a derangement of a strict partition λ , then we define $Q_{\mu}(x) = \text{sgn}(\mu)Q_{\lambda}(x)$, where we have put $\text{sgn}(\mu) = \text{sgn}(\sigma)$.

The Q -functions are related to the power sums via the character values of the symmetric group. An explicit expression is given as follows. Let $\zeta_{\lambda}(\pi)$ ($\lambda \in \mathcal{SP}_n$) be the value of the irreducible projective character ζ_{λ} of the symmetric group $\mathfrak{S}_n$ at the class $\pi = (1^{\pi_1} 3^{\pi_3} 5^{\pi_5} \dots)$ and set $z_{\pi} = \prod_{j \geq 1} \pi_j! j^{\pi_j}$. Then we have

$$Q_{\lambda}(x) = \sum_{\pi} 2^{[(\ell(\lambda) + \ell(\pi) + 1)/2]} z_{\pi}^{-1} \zeta_{\lambda}(\pi) p_1^{\pi_1} p_3^{\pi_3} p_5^{\pi_5} \dots,$$

where the summation runs over all odd partitions $\pi = (1^{\pi_1} 3^{\pi_3} 5^{\pi_5} \dots)$ of size n , and $\ell(\lambda)$ denotes the length of the partition λ .

The following formula is of importance for our purpose.

Theorem 1.1. *Let $j \geq 1$ be odd and $\lambda = (\lambda_1, \dots, \lambda_l) \in \mathcal{SP}$. Then*

$$p_j Q_\lambda(x) = \frac{1}{2} \sum_{i=1}^{(j-1)/2} (-1)^i Q_{(\lambda; j-i, i)}(x) + \frac{1}{2} Q_{(\lambda; j)}(x) + \sum_{i=1}^l Q_{\lambda+j\epsilon_i}(x),$$

where $(\lambda; j-i, i) = (\lambda_1, \dots, \lambda_l, j-i, i)$, $(\lambda; j) = (\lambda_1, \dots, \lambda_l, j)$ and $\epsilon_i = (\delta_{ik})_{1 \leq k \leq l}$.

This is a direct consequence of [J, Theorem 2.4] by looking at the correspondence between removal of *strips/double strips* and that of *j-bars*.

§2 Reduced Q -functions and their linear relations

Let $\Gamma_{\mathbb{Q}}^{(r)} = \mathbb{Q}[p_j; j \geq 1, \text{ odd and } j \not\equiv 0 \pmod{r}]$ and define the r -reduced Q -functions by

$$Q_\lambda^{(r)}(x) = Q_\lambda(x)|_{p_r=p_{3r}=p_{5r}=\dots=0} \in \Gamma_{\mathbb{Q}}^{(r)}.$$

Proposition 2.1. *Let λ be a strict partition and $\lambda^q = (\lambda^0, \dots, \lambda^t)$ ($t = (r-1)/2$) be the r -bar quotient of λ . Then the set $\{Q_\lambda^{(r)}(x); \lambda^0 = \emptyset\}$ gives a basis of $\Gamma_{\mathbb{Q}}^{(r)}$.*

To prove this proposition we need the following lemma.

Lemma 2.2. *Let V be an infinite dimensional $\mathbb{Q}$ -vector space with basis $\{v_\lambda; \lambda \in \mathcal{SP}\}$ and define $v_{\sigma\lambda} = \text{sgn}(\sigma)v_\lambda$ for permutations σ ; $v_\lambda = 0$ if $\lambda \notin \mathcal{SP}$.*

(1) *If we set*

$$V_1 := V \Big/ \sum_{\lambda \in \mathcal{SP}} \mathbb{Q} \left(\sum_{j \geq 1, \text{ odd}} \left(\frac{1}{2} \sum_{i=1}^{(j-1)/2} (-1)^i v_{(\lambda; j-i, i)} + \frac{1}{2} v_{(\lambda; j)} + \sum_{i=1}^{\ell(\lambda)} v_{\lambda+j\epsilon_i} \right) \right),$$

then $V_1 \cong \mathbb{Q}$.

(2) If we set

$$V_r := V \Big/ \sum_{\lambda \in \mathcal{SP}} \mathbb{Q} \left(\sum_{j \geq 1, \text{odd}} \left(\frac{1}{2} \sum_{i=1}^{(rj-1)/2} (-1)^i v_{(\lambda; rj-i, i)} + \frac{1}{2} v_{(\lambda; rj)} + \sum_{i=1}^{\ell(\lambda)} v_{\lambda + rj\epsilon_i} \right) \right),$$

then $V_r \cong \Gamma_{\mathbb{Q}}^{(r)}$.

Proof. (1) There is a canonical linear surjection γ from $\Gamma_{\mathbb{Q}}$ to V_1 which maps $Q_{\lambda}(x)$ to v_{λ} for $\lambda \in \mathcal{SP}$. The kernel of γ coincides with the maximal ideal $I = (p_1, p_3, p_5, \dots)$ because of Theorem 1.1. Since the algebra $\Gamma_{\mathbb{Q}}/I$ is isomorphic to $\mathbb{Q}$, so is V_1 .

(2) Consider a linear surjection $\gamma^{(r)} : \Gamma_{\mathbb{Q}} \longrightarrow V_r$ defined by $\gamma^{(r)}(Q_{\lambda}(x)) = v_{\lambda}$ for $\lambda \in \mathcal{SP}$. If we set $\mathcal{I}^{(r)} = (p_r, p_{2r}, p_{3r}, \dots)$, then, by Theorem 1.1, we see that $\gamma^{(r)}(\mathcal{I}^{(r)}) = 0$ and can define a linear surjection $\bar{\gamma}^{(r)} : \Gamma_{\mathbb{Q}}/\mathcal{I}^{(r)} \longrightarrow V_r$. On the other hand we can define a linear surjection $\eta : V_r \longrightarrow \Gamma_{\mathbb{Q}}^{(r)}$ by $\eta(v_{\lambda}) = Q_{\lambda}^{(r)}(x)$ for $\lambda \in \mathcal{SP}$. The composition $\eta \circ \bar{\gamma}^{(r)} : \Gamma_{\mathbb{Q}}/\mathcal{I}^{(r)} \longrightarrow \Gamma_{\mathbb{Q}}^{(r)}$ gives a linear isomorphism. Hence η is a linear isomorphism as desired. ■

Proof of Proposition 2.1. We give a proof only for the case $r = 3$ in order to avoid the too complicated notation. The general case can be shown in a same fashion.

First we shall see that the set $\{Q_{\lambda}^{(3)}; \lambda^0 = \emptyset\}$ spans $\Gamma_{\mathbb{Q}}^{(3)}$. We introduce a filtration on $\Gamma_{\mathbb{Q}}^{(3)}$ by $F_n \Gamma_{\mathbb{Q}}^{(3)} = \sum_{\lambda; |\lambda^0| \leq n} \mathbb{Q} Q_{(\lambda^c, \lambda^0, \lambda^1)}^{(3)}(x)$ and the associated graded module $\overline{F}(\Gamma_{\mathbb{Q}}^{(3)}) :=$

$\bigoplus_{n \geq 0} F_n \Gamma_{\mathbb{Q}}^{(3)} / F_{n-1} \Gamma_{\mathbb{Q}}^{(3)}$, where $F_{-1} \Gamma_{\mathbb{Q}}^{(3)} = \{0\}$. For any positive odd integer j the 3-reduced

Q -functions $Q_{\lambda}^{(3)}(x)$ satisfy

$$\begin{aligned} 0 &= \frac{1}{2} \sum_{i=1}^{(3j-1)/2} Q_{(\lambda; 3j-i, i)}^{(3)}(x) + \frac{1}{2} Q_{(\lambda; 3j)}^{(3)}(x) + \sum_{i=1}^{\ell(\lambda)} Q_{\lambda + 3j\epsilon_i}^{(3)}(x) \\ &= \frac{1}{2} \sum_{i=1}^{(j-1)/2} Q_{(\lambda^c, (\lambda^0; j-i, i), \lambda^1)}^{(3)}(x) + \frac{1}{2} Q_{(\lambda^c, (\lambda^0; j), \lambda^1)}^{(3)}(x) + \sum_{i=1}^{\ell(\lambda^0)} Q_{(\lambda^c, \lambda^0 + j\epsilon_i, \lambda^1)}^{(3)}(x) \end{aligned}$$

$$+ \frac{1}{2} \sum_{i \not\equiv 0 \pmod{3}} Q_{(\lambda^c, \lambda^0, (\lambda; 3j-i, i)^1)}^{(3)} + \sum_{i=1}^{\ell(\lambda^1)} Q_{(\lambda^c, \lambda^0, \lambda^1 + j\epsilon_i)}^{(3)}(x).$$

If we write $\overline{Q}_\lambda^{(3)}(x) = Q_\lambda^{(3)}(x) + F_{n-1}\Gamma_{\mathbb{Q}}^{(3)}$ for $|\lambda^0| = n$, then we see that in $\overline{F}(\Gamma_{\mathbb{Q}}^{(3)})$

$$\sum_{i=1}^{\ell(\lambda^0)} \overline{Q}_{(\lambda^c, \lambda^0 + j\epsilon_i, \lambda^1)}^{(3)}(x) + \frac{1}{2} \overline{Q}_{(\lambda^c, (\lambda^0; j), \lambda^1)}^{(3)}(x) + \frac{1}{2} \sum_{i=1}^{(j-1)/2} \overline{Q}_{(\lambda^c, (\lambda^0; j-i, i), \lambda^1)}^{(3)}(x) = 0.$$

By applying Lemma 2.2 (1) it turns out that $\sum_{\lambda^0 \in \mathcal{SP}} \mathbb{Q} \overline{Q}_{(\lambda^c, \lambda^0, \lambda^1)}^{(3)}(x) = \mathbb{Q} \overline{Q}_{(\lambda^c, \emptyset, \lambda^1)}^{(3)}(x)$. This

proves that $\{Q_\lambda^{(3)}(x); \lambda^0 = \emptyset\}$ spans $\Gamma_{\mathbb{Q}}^{(3)}$.

By looking at the one-to-one correspondence between the strict partitions and the tuples of r -cores and r -quotients, we see that the number of those strict partitions $\lambda \in \mathcal{SP}_n$ such that $\lambda^0 = \emptyset$ coincides with the dimension of the subspace of homogeneous polynomials of degree n in $\Gamma_{\mathbb{Q}}^{(r)}$, where we count $\deg p_j = j$. This completes the proof. $\blacksquare$

We shall state the linear relations satisfied by the r -reduced Q -functions. To this end, we define two types of Littlewood-Richardson-like coefficients $f_{\lambda_1 \dots \lambda_t}^\nu$ ($\lambda_1, \dots, \lambda_t, \nu \in \mathcal{SP}$) and $h_{\lambda\mu}^\nu$ ($\lambda \in \mathcal{SP}$, $\mu, \nu \in \mathcal{P}$), respectively by

$$P_{\lambda_1}(x) \cdots P_{\lambda_t}(x) = \sum_{\nu \in \mathcal{SP}} f_{\lambda_1 \dots \lambda_t}^\nu P_\nu(x),$$

$$P_\lambda(x) s_\mu(x) = \sum_{\nu \in \mathcal{P}} h_{\lambda\mu}^\nu s_\nu(x),$$

where we set $P_\lambda(x) = 2^{-\ell(\lambda)} Q_\lambda(x)$ for $\lambda \in \mathcal{SP}$. Note that $f_{\lambda_1 \dots \lambda_t}^\nu$ and $h_{\lambda\mu}^\nu$ are non-negative integers.

Theorem 2.3. *Let λ be a strict partition, λ^c and $\lambda^q = (\lambda^0, \lambda^1, \dots, \lambda^t)$ be its r -bar core and r -bar quotient, respectively. Then*

$$Q_\lambda^{(r)}(x) = 2^{[(\ell(\lambda) + \ell(\lambda^0))/2]} (-1)^{|\lambda^0|} \delta_{\bar{r}}(\lambda) \sum_{\mu, \nu_1, \dots, \nu_t} 2^{-[\ell(\mu)/2]} \delta_{\bar{r}}(\mu) f_{\nu_1 \dots \nu_t}^{\lambda^0} h_{\nu_1 \lambda^1}^{\mu^1} \cdots h_{\nu_t \lambda^t}^{\mu^t} Q_\mu^{(r)}(x),$$

where summation runs over the strict partitions $\nu_1, \dots, \nu_t$ and μ such that $|\mu| = |\lambda|$, $\mu^c = \lambda^c$ and $\mu^0 = \emptyset$.

Proof. Again we give a proof only for the case $r = 3$. For the general case we should notice the following identity :

$$Q_\nu(x^{(1)}, x^{(2)}, \dots, x^{(t)}) = \sum_{\lambda_1, \dots, \lambda_t} f_{\lambda_1 \dots \lambda_t}^\nu Q_{\lambda_1}(x^{(1)}) Q_{\lambda_2}(x^{(2)}) \cdots Q_{\lambda_t}(x^{(t)}),$$

where summation runs over the strict partitions $\lambda_1, \dots, \lambda_t$. Here $(x^{(1)}, x^{(2)}, \dots, x^{(t)})$ are t sets of variables and $Q_\nu(x^{(1)}, \dots, x^{(t)})$ denotes the Q -function in the set of variables $(x_1^{(1)}, x_2^{(1)}, \dots, x_1^{(2)}, x_2^{(2)}, \dots, x_1^{(t)}, x_2^{(t)}, \dots)$.

Set

$$F_\lambda(x) = \text{sgn}(\lambda) 2^{[(\ell(\lambda) + \ell(\lambda^0))/2]} (-1)^{|\lambda^0|} \delta_{\bar{3}}(\lambda) \sum_{\mu} 2^{-[\ell(\mu)/2]} \delta_{\bar{3}}(\mu) h_{\lambda^0 \lambda^1}^{\mu^1} Q_\mu^{(3)}(x).$$

We shall prove that

$$\sum_{i=1}^{(3j-1)/2} (-1)^i F_{(\lambda; 3j-i, i)}(x) + \frac{1}{2} F_{(\lambda; 3j)}(x) + \sum_{i=1}^{\ell(\lambda)} F_{\lambda+3j\epsilon_i}(x) = 0$$

for any positive odd integer j . For our purpose it is enough to show that the coefficient of $Q_\mu^{(3)}(x)$ vanishes for each $\mu \in \mathcal{SP}$. Since corollary in [MO, (3.8)] leads to

$$\begin{aligned} & \delta_{\bar{3}}(\tilde{\lambda}, \lambda) \\ &= \begin{cases} \text{sgn}(\tilde{\lambda}) \text{sgn}(\tilde{\lambda}^k) & \text{for } \tilde{\lambda} = \lambda + 3j\epsilon_i \text{ } (k = 0, 1, \text{ } i = 1, \dots, \ell(\lambda)) \\ \text{sgn}(\tilde{\lambda}) \text{sgn}(\tilde{\lambda}^0) & \text{for } \tilde{\lambda} = (\lambda; 3j) \text{ and } (\lambda; 3j - 3j', 3j') \text{ } (j' \geq \frac{j-1}{2}) \\ (-1)^i \text{sgn}(\tilde{\lambda}) \text{sgn}(\tilde{\lambda}^1) & \text{for } \tilde{\lambda} = (\lambda; 3j - i, i) \text{ } (i \geq \frac{3j-1}{2}, i \not\equiv 0 \pmod{r}) \end{cases}, \end{aligned}$$

our equation reads

$$\frac{1}{2} \sum_{i < j/2} 2^2 (-1)^i \text{sgn}((\lambda^0; j - i, i)) (-1)^j h_{(\lambda^0; j-i, i)\lambda^1}^{\mu^1}$$

$$\begin{aligned}
& + \frac{1}{2} \sum_{i \not\equiv 0 \pmod{3}} 2 \operatorname{sgn}((\lambda; 3j - i, i)^1) h_{\lambda^0(\lambda; 3j - i, i)^1}^{\mu^1} + \frac{1}{2} \operatorname{sgn}((\lambda^0; j)) 2(-1)^j h_{(\lambda^0; j)\lambda^1}^{\mu^1} \\
& + \sum_i \operatorname{sgn}((\lambda + 3j\epsilon_i)^0) (-1)^j h_{(\lambda + 3j\epsilon_i)^0\lambda^1}^{\mu^1} + \sum_i \operatorname{sgn}((\lambda + 3j\epsilon_i)^1) h_{\lambda^0(\lambda + 3j\epsilon_i)^1}^{\mu^1} \\
& = 0.
\end{aligned}$$

Multiplying both sides by $s_{\mu^1}(x)$ and taking a summation over μ^1 , we see that the equation reduces to

$$\begin{aligned}
& (-1)^j \left\{ \frac{1}{2} \sum_{i < j/2} (-1)^i Q_{(\lambda^0; j - i, i)}(x) s_{\lambda^1}(x) + \frac{1}{2} Q_{(\lambda^0; j)}(x) s_{\lambda^1}(x) + \sum_i Q_{\lambda^0 + j\epsilon_i}(x) s_{\lambda^1}(x) \right\} \\
& + \left\{ \sum_i Q_{\lambda^0}(x) s_{\lambda^1 + j\epsilon_i}(x) + \sum_{\substack{i < j/2 \\ i \not\equiv 0 \pmod{3}}} Q_{\lambda^0}(x) s_{(\lambda, 3j - i, i)^1}(x) \right\} = 0.
\end{aligned}$$

Note that there is a one-to-one correspondence between the $3j$ -bars in λ and the j -bars in $\lambda^q = (\lambda^0, \lambda^1)$ (a j -bar in λ^q means a j -bar in λ^0 or a j -hook in λ^1). Hence we have

$$-(p_j Q_{\lambda^0}(x)) s_{\lambda^1}(x) + Q_{\lambda^0}(x) (p_j s_{\lambda^1}(x)) = 0.$$

By Lemma 2.2 (3) there exists a linear isomorphism sending $F_\lambda(x)$ to $Q_\lambda^{(3)}(x)$ and it turns out to be the identity since $Q_\lambda^{(3)}(x)$ equals $F_\lambda(x)$ for the basis given in Proposition 2.1. ■

References

- [ANY1] S. Ariki, T. Nakajima and H.-F. Yamada, 'Weight vectors of the basic $A_1^{(1)}$ -module and the Littlewood-Richardson rule', *J. Phys. A: Math. Gen.* 28 (1995) L357–L361.
- [ANY2] S. Ariki, T. Nakajima and H.-F. Yamada, 'Reduced Schur functions and the Littlewood-Richardson coefficients', *Preprint*.

- [J] T. Józefiak, 'Schur Q -functions and applications', Proc. Hyderabad Conf. on Algebraic Groups, S. Ramanan ed. Manoj Prakashan (1991) 205–224.
- [M] I. G. Macdonald 'Symmetric Functions and Hall Polynomials, 2nd ed.' Oxford (1995).
- [MO] A. O. Morris and J. B. Olsson, 'On p -quotients for spin characters', *J. Algebra* 119 (1988) 51–82.
- [NY] T. Nakajima and H.-F. Yamada, 'Basic representations of $A_{2l}^{(2)}$ and $D_{l+1}^{(2)}$ and the polynomial solutions to the reduced BKP hierarchies', *J. Phys. A: Math. Gen.* 27 (1994) L171–L176.
- [O] J. B. Olsson 'Combinatorics and Representations of Finite Groups' Lecture Notes, University of Essen, vol. 20 (1993).